

Newborn Rotation Checklist

Topic	Complete
Counsel at least 3 mothers on post-partum on breastfeeding	
Counsel at least 3 mothers in an outpatient setting on breastfeeding	
Evaluate at least 5 infants for weight check	
Treat at least 2 infants with phototherapy	
Treat at least 2 infants with antibiotics for any indication	
Correctly identify at least one infant with caput succedaneum	
Correctly identify at least one infant with cephalohematoma and counsel parent on prognosis	
Demonstrate one newborn exam to an attending	
Describe the murmur and determine the best course of follow-up for at least two infants with heart murmurs	
Identify at least one infant with the following: milia, nevus flammeus, pustular melanosis, miliaria, infantile acne, erythema toxicum	
Administer IV fluids to at least 2 infants	
Manage at least 2 infants as they advance to nipple feeds	
Manage at least 2 infants of diabetic mothers	
Manage at least 1 infant with respiratory distress	
Evaluate at least 1 infant with jitteriness	
Resuscitate at least 1 infant who requires PPV or, if no infant requires this during the rotation, describe when PPV is indicated and demonstrate PPV technique on a dummy.	
Insert at least one umbilical catheter (on an infant if possible, if not, utilizing a length of umbilical cord)	
Manage either one infant with neonatal abstinence syndrome, an infant born to a mother with a history of substance abuse, or an infant with a positive urine toxicology screen	
Review workup of feeding intolerance, vomiting, abdominal distention, and no stool within the first 48 hours of life with an attending	
Number of C-Sections - Please add a hash mark for each C-Section:	